

URUGUAY

Distribución de casos y tasas específicas por intervalo etario y sexo


Todos los sitios reunidos Bienio 1996 - 1997

Edades	Hombres		Mujeres	
	Nº	Tasa Esp.	Nº	Tasa Esp.
0-4	14	5,14	17	6,41
5-9	20	7,42	23	8,80
10-14	26	9,89	22	8,53
15-19	49	18,31	36	13,93
20-24	77	31,17	74	30,41
25-29	73	33,84	128	58,50
30-34	125	58,10	248	111,79
35-39	159	76,63	367	168,86
40-44	306	163,06	603	304,87
45-49	460	274,52	734	413,43
50-54	738	484,16	887	544,16
55-59	1.199	861,32	1.025	661,75
60-64	1.741	1.300,40	1.269	820,03
65-69	2.223	1.812,20	1.567	1.039,00
70-74	2.228	2.432,30	1.545	1.275,70
75-79	1.884	3.131,60	1.467	1.605,70
80-84	1.196	3.347,20	1.243	1.894,30
85 y más	965	4.035,80	1.198	2.268,00
Sin edad	23		29	
Total	13.506		12.482	
Tasa aj.		325,35		250,75
Tasa cr.		439,54		381,26

Distribución porcentual de la incidencia Localizaciones más frecuentes Bienio 1996 - 1997

HOMBRES

		Casos	%
1	PULMON	2.716	20,11
2	PROSTATA	1.985	14,70
3	COLO-RECTO	1.437	10,64
4	PIEL	962	7,12
5	ESTOMAGO	804	5,95
	PRIM. DESC.	831	6,15



MUJERES

		Casos	%
1	MAMA	3.460	27,72
2	COLO-RECTO	157	12,59
3	CUELLO DE UTERO	978	7,84
4	PIEL	781	6,26
5	PULMON	515	4,13
	PRIM. DESC.	698	5,59

