



MONTEVIDEO

Distribución de casos y tasas específicas por intervalo etario y sexo


Todos los sitios reunidos
Bienio 1996 - 1997

Edades	Hombres		Mujeres	
	Nº	Tasa Esp.	Nº	Tasa Esp.
0-4	5	5,04	11	11,54
5-9	8	7,88	9	9,22
10-14	11	11,25	11	11,40
15-19	20	18,17	12	11,07
20-24	34	30,12	37	32,40
25-29	37	39,35	59	60,87
30-34	75	82,55	105	109,96
35-39	79	90,53	164	169,94
40-44	150	193,38	270	307,89
45-49	231	345,52	350	445,09
50-54	351	569,79	418	555,04
55-59	575	1.020,30	527	726,37
60-64	885	1.570,60	670	889,73
65-69	1.142	2.127,70	918	1.213,00
70-74	1.148	2.860,00	872	1.408,40
75-79	988	3.881,50	820	1.783,60
80-84	594	3.936,90	675	1.981,40
85 y más	459	4.541,40	683	2.455,70
Sin edad	10		24	
Total	6.802		6.635	
Tasa aj.		386,68		270,74
Tasa cr.		541,20		462,10

Distribución porcentual de la incidencia Localizaciones más frecuentes Bienio 1996 - 1997

HOMBRES

		Casos	%
1	PULMON	1.352	19,88
2	PROSTATA	1.043	15,33
3	COLO-RECTO	750	11,03
4	PIEL	473	6,95
5	VEJIGA	470	6,91
	PRIM. DESC.	371	5,45



MUJERES

		Casos	%
1	MAMA	2.049	30,88
2	COLO-RECTO	867	13,07
3	PIEL	397	5,98
4	CUELLO DE UTERO	318	4,79
5	PULMON	284	4,28
	PRIM. DESC.	369	5,56

