



INTERIOR

Distribución de casos y tasas específicas por intervalo etario y sexo


Todos los sitios reunidos Bienio 1996 - 1997

Edades	Hombres		Mujeres	
	Nº	Tasa Esp.	Nº	Tasa Esp.
0-4	9	5,19	6	3,53
5-9	12	7,14	14	8,55
10-14	15	9,08	11	6,81
15-19	29	18,41	24	16,00
20-24	43	32,05	37	28,64
25-29	36	29,59	69	56,62
30-34	50	40,23	143	113,16
35-39	80	66,54	203	168,00
40-44	156	141,70	333	302,47
45-49	229	227,39	384	388,26
50-54	387	426,08	469	534,81
55-59	624	753,17	498	604,81
60-64	856	1.103,90	599	753,96
65-69	1.081	1.566,70	649	863,71
70-74	1.080	2.098,60	673	1137,00
75-79	896	2.581,70	647	1425,50
80-84	602	2.916,20	568	1800,20
85 y más	506	3.665,60	515	2059,20
Sin edad	13		5	
Total	6.704		5.847	
Tasa aj.		280,58		231,76
Tasa cr.		369,17		318,11

Distribución porcentual de la incidencia Localizaciones más frecuentes Bienio 1996 - 1997

HOMBRES

		Casos	%
1	PULMON	1364	20,35
2	PROSTATA	942	14,05
3	COLO-RECTO	687	10,25
4	PIEL	489	7,29
5	ESTOMAGO	485	7,23
	PRIM. DESC.	460	6,86



MUJERES

		Casos	%
1	MAMA	1411	24,13
2	COLO-RECTO	705	12,06
3	CUELLO UTERINO	660	11,29
4	PIEL	384	6,57
5	ESTOMAGO	261	4,46
	PRIM. DESC.	329	5,63

